



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....



**Group A Streptococcus – Infection Prevention and Control**  
**Patients Name:** .....  
**Patient CHI:** .....  
**GAS Isolated on (Date):** .....

The patient should be isolated in a single room with standard and transmission based precautions, until they have received 48 hours of appropriate antibiotics **and** there is significant clinical improvement. The following should be implemented:

- The patient room/bed space and equipment must be cleaned twice per day with chlorine based detergent.
- A yellow disposable apron and gloves must be worn for all direct contact with the patient or the patient’s environment.
- A fluid resistant surgical mask should be worn if there is a risk of blood or body fluid splash.
- An FFP3 mask, yellow disposable apron, gloves and eye protection must be worn for all aerosol generating procedures.
- A yellow IPC Isolation sign must be on the door to the room and the door must be closed.
- The Group A Streptococcus care checklist should be completed daily in conjunction with the NHS GGC GAS SOP.

**This advice has been given by:** ..... (IPCN)  
**Date:** .....