



**Infection Prevention and Control Advice (Influenza)**

Patient Name & CHI.....

**Influenza A / B isolated :** .....

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent .
- Disposable yellow apron, gloves and surgical mask should be worn by HCW when delivering direct care / within 1m of the patient.
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn.
- The door to the room must be closed.
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene.
- The IPC influenza virus care checklist should be completed daily, including risk assessment when TBPs cannot be implemented.

IPCN: .....

Date: .../.../.....



**Infection Prevention and Control Advice (Influenza)**

Patient Name & CHI.....

**Influenza A / B isolated :** .....

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent .
- Disposable yellow apron, gloves and surgical mask should be worn by HCW when delivering direct care / within 1m of the patient.
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn.
- The door to the room must be closed.
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene.
- The IPC influenza virus care checklist should be completed daily, including risk assessment when TBPs cannot be implemented.

IPCN: .....

Date: .../.../.....



**Infection Prevention and Control Advice (Influenza)**

Patient Name & CHI.....

**Influenza A / B isolated :** .....

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent .
- Disposable yellow apron, gloves and surgical mask should be worn by HCW when delivering direct care / within 1m of the patient.
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn.
- The door to the room must be closed.
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene.
- The IPC influenza virus care checklist should be completed daily, including risk assessment when TBPs cannot be implemented.

IPCN: .....

Date: .../.../.....



**Infection Prevention and Control Advice (Influenza)**

Patient Name & CHI.....

**Influenza A / B isolated :** .....

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent .
- Disposable yellow apron, gloves and surgical mask should be worn by HCW when delivering direct care / within 1m of the patient.
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn.
- The door to the room must be closed.
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene.
- The IPC influenza virus care checklist should be completed daily, including risk assessment when TBPs cannot be implemented.

IPCN: .....

Date: .../.../.....