



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....



**Influenza - Infection Prevention and Control**

**Patient Name:** .....  
**Patient CHI:** .....

**Influenza A / B Date Isolated:** ---/---/---

The patient should be isolated in a single room with standard and transmission based precautions, until at least 24 hours free of coryzal symptoms OR patient has completed a course of antivirals. The following must be implemented:

- The patient room /bed space and equipment must be cleaned twice per day with chlorine based detergent
- Disposable yellow apron, gloves and surgical mask must be worn by HCW when delivering direct care / within 1m of the patient
- During aerosol generating procedures, an FFP3 mask and eye protection must be worn
- The door to the room must be closed
- Supply patient with disposable tissues and encourage good cough and respiratory hygiene
- The IPC influenza virus care checklist must be completed daily, including risk assessment if TBPs cannot be implemented.

**IPCN:** ..... **Date:** .../.../.....