

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	1 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

SOP Objective

To provide Healthcare Workers (HCW) with details of the care required to prevent cross-infection in children with *Clostridioides difficile* Infection (CDI).

This SOP applies to all staff employed by NHS Greater Glasgow & Clyde and locum staff on fixed term contracts and volunteer staff.

KEY CHANGES FROM THE PREVIOUS VERSION OF THIS POLICY

Document Control Summary

Approved by and date	Board Infection Control Committee 15 th June 2020
Date of Publication	17 th June 2020
Developed by	Infection Control Policy Sub-Group
Related Documents	National Infection Prevention and Control Manual NHSGGC SOP CDI (Adults) NHSGGC SOP Hand Hygiene NHSGGC SOP Outbreak NHSGGC SOP Cleaning of Near Patient Equipment NHSGGC SOP Terminal Clean of Isolation Rooms NHSGGC SOP Twice daily Clean of Isolation Rooms Antimicrobial Prescribing Policies
Distribution/ Availability	NHSGGC Infection Prevention and Control Policy Manual and the Internet http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control
Lead Manager	Board Infection Control Manager
Responsible Director	Board Medical Director

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>


	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	2 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

CONTENTS

CDI Paediatric Aide Memoire.....	3
1. Responsibilities.....	4
2. General Information on <i>Clostridioides difficile</i> Infection (CDI)	6
3. Transmission Based Precautions for CDI	8
4. Evidence Base	16
Appendix 2 – Bowel Movement (adapted from the Bristol Stool Scale).....	17

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	3 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

CDI Paediatric Aide Memoire



	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	4 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

1. Responsibilities

Healthcare Workers (HCWs) must:

- Follow this SOP.
Commence a CDI Care Checklist while patient is symptomatic, update daily and complete the risk assessment for any aspect of transmission based precautions (TBP) for CDI that cannot be implemented

[Clostridioides Difficile – IPC Care checklist](#)

- Inform their line manager and a member of the Infection Prevention and Control Team if this SOP cannot be followed.
- Provide written and verbal information on CDI for patients and their relatives as appropriate

[Clostridioides Difficile Fact Sheet](#)

Senior Charge Nurse (SCN) must:

- Ensure that the IPC Care checklist is in place while patient is deemed infectious.
- Ensure that written information is provided / available for patients and relatives.
- Ensure a failure to isolate risk assessment is in place if any aspect of TBPs for CDI cannot be implemented

Managers must:

- Support HCWs and IPCTs in following this SOP.
- Cascade new SOPs to clinical staff after approval by the Board Infection Control Committee (BICC).

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	5 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Infection Prevention and Control Teams (IPCTs) must:

- Keep this SOP up-to-date.
- Provide education opportunities on this SOP.
- Monitor epidemiology of *Clostridioides difficile* Infection (CDI) within healthcare facility(ies) and advise on infection prevention and control precautions as necessary.
- Advise and support HCWs to undertake a Risk Assessment if unable to follow this SOP.

Occupational Health Service (OHS) must:

- Advise HCW regarding possible infection exposure and return to work issues as necessary

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	6 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

2. General Information on *Clostridioides difficile* Infection (CDI)

<i>Communicable Disease/ Alert Organism</i>	<p><i>Clostridioides difficile</i> is a Gram positive, anaerobic, spore forming, toxin producing gastrointestinal bacillus. However, recent studies have shown that <i>C. difficile</i> is an emerging pathogen in the paediatric setting, causing a range of illness; from mild diarrhoea to life changing conditions such as pseudo-membranous colitis, toxic megacolon, intestinal perforation and septic shock. It is imperative that clinical judgement is exercised in order that aetiologies are appropriately investigated.</p>
<i>Case Definition</i>	<p>A child (3-16 years of age) has a diagnosis of CDI if they have a stool specimen positive for CD toxin, diarrhoea (Bristol stool chart <u>5-7</u>) and one or more of the following:</p> <ul style="list-style-type: none"> • Significant co-morbidities i.e. haematology/oncology ; gastrointestinal • Severe GI disease with bloody diarrhoea and an unlikely alternative diagnosis • Strong clinical suspicion • Antibiotic therapy in the last 4 weeks (especially ciprofloxacin)
<i>Case Definition : Determination of source</i>	<p>Hospital acquired CDI is defined as when a patient has had onset of symptoms at least 48 hours following admission to a hospital</p> <p>Healthcare associated CDI is defined as when a patient has had onset of symptoms up to four weeks after discharge from a hospital</p> <p>Indeterminate cases of CDI is defined as a patient who was discharged from a hospital 4–12 weeks before the onset of symptoms.</p> <p>Community associated CDI Is defined as a patient with onset of symptoms while outside a hospital and without discharge from a hospital within the previous 12 weeks – or with onset</p>

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	7 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

	of symptoms within 48 hours following admission to a hospital without stay in a hospital within the previous 12 weeks
<i>Mode of Spread</i>	There is evidence of both direct and indirect spread through the hands of HCWs and patients; and environmental contamination via equipment and instruments, e.g. commodes, bedpans and washbowls. <i>C. difficile</i> produces spores which can survive for long periods in the environment. Environmental cleaning is paramount.
<i>Incubation Period</i>	Potentially up to 12 weeks.

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	8 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

3. Transmission Based Precautions for CDI

<i>Accommodation (Patient Placement)</i>	<p>The patient should be placed in a single room, preferably with ensuite or own commode. The door to the room should be closed when not in use and a yellow IPCT sign placed on the door. If a side room is unavailable the IPCT will help the clinical team to undertake a risk assessment and advise where to nurse the patient.</p> <p>Precautions should continue until the patient has been asymptomatic for 48 hours and bowel movements have returned to normal or, on the advice of a member of the IPCT.</p>
<i>Care Checklist available</i>	Yes. CDI Care Checklist
<i>Clinical/ Healthcare Waste</i>	All non-sharps waste should be designated as Healthcare/Clinical Waste (HCW) and placed in an orange clinical waste bag within the room. Please refer to the NHSGCC Waste Management Policy .
<i>Contacts</i>	Specimens should not be sent from patients deemed to be contacts unless they develop loose stools, where there is no other cause for this.
<i>Domestic Services/ Facilities</i>	Domestic staff must follow the NHSGGC SOP for Twice Daily Clean of Isolation Rooms . Cleans should be undertaken at least four hours apart

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	9 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Equipment	<p>Patient equipment, e.g. commode, BP cuff, washbowl should be allocated to the patient until no longer considered infectious. Consider single-use or single patient use equipment. Commodes should be decontaminated after each use with chlorine based detergent.</p> <p>Twice Daily Clean of Isolation Rooms SOP</p>
Hand Hygiene	<p>Alcohol gel hand rub and chlorhexidine are <u>not</u> effective against CDI: Soap and water must be used for all patients with loose stools.</p> <p>Hand hygiene is the single most important measure to prevent cross infection with CDI. Hands must be decontaminated before and after each direct patient contact, after contact with the environment, after exposure to body fluids and before any aseptic tasks. Patients should be encouraged to carry out thorough hand hygiene. If a child is unable to decontaminate their hands then hand hygiene should be carried out by a HCW or patient carer.</p> <p>Please refer to NHSGGC Hand Hygiene SOP</p> <p>Visitors should also be instructed to wash their hands with soap and water</p>
Health Protection Scotland (HPS) Trigger Tool	<p>The Health Protection Scotland (HPS) Trigger Tool must be completed by the IPCT and Clinical Staff if there are two or more HAI CDI cases in the same ward in a two week period. IPCNs and ward staff will complete the trigger daily until the trigger is no longer in place i.e. one or both patients are no longer symptomatic or have been discharged. The following actions will be taken by the IPCT when a trigger is met:</p> <ul style="list-style-type: none"> - Request a terminal clean of the ward at the start of the trigger - Advise on enhanced IPC precautions to be in place. - Undertake IPC audit (if not done in last 3 months) hand hygiene audit - Ask the antimicrobial pharmacist to review prescribing

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	10 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

	<p>Findings will be reported to the SCN and ward staff who should liaise with IPC and pharmacy colleagues on any actions required as a result.</p> <p>Following this, should another case of HAI CDI emerge, the IPCT will complete a PAG to determine the requirement for an IMT and ward closure.</p>
<i>Linen</i>	<p>Treat used linen as soiled/ infected, i.e. place in a water soluble alginate bag then a clear bag tied and then into a laundry bag. (Brown polythene bag used in Mental Health areas)</p> <p>Please refer to National Guidance on the safe management of linen.</p>
<i>Moving between wards, hospitals and departments (including theatres)</i>	<p>Except in clinical emergencies, transfer of patients who have not been symptom-free and passed normal stool for 48-hours is not advisable.</p> <p>However, acute receiving units have a high patient turnover and transfer of patients is necessary for effective patient flow and to ensure that patients receive the appropriate care within their specialty. Therefore, Receiving areas MUST be informed of the patient's condition before the patient is transferred and the requirement for a single room.</p> <p>Please follow NHSGGC SOP Terminal Clean of Isolation Rooms.</p>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	11 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Notice for Door	<p>The yellow IPC isolation sign must be placed on the door to the patient's room.</p> <p>In Mental Health Services (MHS), on advice of IPCT.</p>
Patient Clothing	<p>Whilst patients are very symptomatic they should be advised to wear hospital gowns.</p> <p>If relatives or carers wish to take personal clothing home, staff must place soiled clothing into a domestic alginate bag and staff must ensure that a Home Laundry Information Leaflet is issued.</p> <p>NB: It should be recorded in the nursing notes that both the advice and information leaflet has been issued.</p>
Patient Information	<p>Inform the patient and / or if relevant, the patient's relative/ carer of their condition and the necessary precautions if required. Answer any questions and concerns they may have. A CDI Fact sheet for patients and their relatives is available to download from the IPCT website.</p> <p>NB: It should be recorded in the nursing notes or Care Checklist that the fact sheet has been issued. IPCTs are available to speak to patients or relatives/ carers if required.</p> <p>CDI Fact Sheet</p>
Personal Protective Equipment (PPE)	<p>To prevent spread through direct contact PPE (disposable gloves and yellow apron) must be worn for all direct contact with the patient or the patient's environment/equipment.</p> <p>If there is a risk of splashing of blood/body fluids, then facial protection i.e. mask/visor should also be considered. Hand hygiene must be performed using liquid soap and water before donning and after removing PPE. Alcohol based hand rub is not effective against CDI.</p>
Precautions required until	<p>Precautions should continue until the patient has been asymptomatic for 48 hours and bowel movements have returned to normal or, on advice of a member of the IPCT.</p> <p>If symptoms recur, reinstate precautions immediately, send further specimens and inform a member of the IPCT.</p>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	12 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Daily and weekly check by IPCT	IPCNs will check daily (Monday -Friday) on the condition of patients with CDI until TBPs are no longer required and thereafter weekly for 4 weeks.																																				
Daily assessment of severity by clinical team	<p>If the patient is confirmed as CDI, and while the patient is symptomatic of loose stools, medical staff are required to undertake a daily severity assessment using the assessment tool below. Daily severity assessments should continue until patient has been asymptomatic for more than 48 hours. Medical staff should consider the need to take bloods to complete the severity score.</p> <p>Severity assessment in paediatric population (3-16 years)</p> <table border="1" data-bbox="584 958 1385 1330"> <thead> <tr> <th>Criteria</th> <th>Yes</th> <th>No</th> <th>Score if Yes</th> </tr> </thead> <tbody> <tr> <td>Diarrhoea >5 times per day</td> <td></td> <td></td> <td>1</td> </tr> <tr> <td>Abdominal pain and discomfort</td> <td></td> <td></td> <td>1</td> </tr> <tr> <td>Rising white cell count</td> <td></td> <td></td> <td>1</td> </tr> <tr> <td>Raised C-reactive protein</td> <td></td> <td></td> <td>1</td> </tr> <tr> <td>Pyrexia >38 °C</td> <td></td> <td></td> <td>1</td> </tr> <tr> <td>Evidence of pseudo membranous colitis</td> <td></td> <td></td> <td>2</td> </tr> <tr> <td>Intensive care unit requirement</td> <td></td> <td></td> <td>2</td> </tr> <tr> <td>Total score</td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p>≥ 5 = severe disease</p> <p>If a patient is assessed as severe the IPCT will:</p> <ul style="list-style-type: none"> • refer to the CDI treatment algorithm (paeds) • Communicate severe cases to the Senior Management Team/ Microbiology and Clinical Teams • IPCT will generate a datix 	Criteria	Yes	No	Score if Yes	Diarrhoea >5 times per day			1	Abdominal pain and discomfort			1	Rising white cell count			1	Raised C-reactive protein			1	Pyrexia >38 °C			1	Evidence of pseudo membranous colitis			2	Intensive care unit requirement			2	Total score			
Criteria	Yes	No	Score if Yes																																		
Diarrhoea >5 times per day			1																																		
Abdominal pain and discomfort			1																																		
Rising white cell count			1																																		
Raised C-reactive protein			1																																		
Pyrexia >38 °C			1																																		
Evidence of pseudo membranous colitis			2																																		
Intensive care unit requirement			2																																		
Total score																																					
Clinical review assessment (CRA) and Reporting of Severe Cases of CDI	A Clinical Review is required if the patient: <ul style="list-style-type: none"> • Has severe or life threatening CDI • Was admitted to ITU for treatment of CDI or its complications • Had endoscopic diagnosis of pseudomembranous colitis with or without toxin confirmation • Had surgery for the complications of CDI (toxic megacolon, perforation or refractory colitis) • Died within 30 days following a diagnosis of CDI where it 																																				

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	13 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

	<p>is recorded as either the primary or a major contributory factor on the death certificate</p> <ul style="list-style-type: none"> • Had persisting CDI where the patient has remained symptomatic and toxin positive despite two courses of appropriate therapy
<i>Deaths due to CDI (Underlying or Contributing)</i>	<p>If death certificate records are not available, the lead IPCN will contact the General Manager (GM) for the service, and advise them that the records are not available. The Lead Infection Prevention and Control Doctor (LIPCD), Infection Prevention and Control Manager (IPCM), Associate Nurse Director, Infection Prevention and Control (ANDIPC), Clinical Services Manager (CSM) and Lead Nurse for the area must be informed of all patients who died in hospital who are or who have been positive for CDI during their current admission, and the cause of death if available. If a Datix has not been completed for a severe case one should be completed at this point.</p> <p>Medical staff completing a death certificate in which CDI is noted (part 1 or 2) should discuss this with the consultant in charge of the patient's clinical care and refer case to the Procurator Fiscals Office. If CDI is placed on part 1, medical staff should inform the CSM and GM for the area.</p> <p>Medical staff should familiarise themselves with NHSGGC Guidance on the Completion of Medical Certificates of Cause of Death.</p>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	14 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Treatment	<p><u>Mild disease:</u> No need to treat if symptoms settle within 24 hours but consider oral metronidazole for 10–14 days if symptoms persist beyond 24 hours.</p> <p><u>Moderate disease:</u> Oral metronidazole for 10–14 days and consider escalation by changing to oral vancomycin if non-resolution of symptoms or decline in severity score.</p> <p><u>Severe disease:</u> Oral vancomycin and IV metronidazole. Colectomy should be considered if there is evidence of caecal dilatation. Surgical review should be considered/discussed with microbiology The Management of Suspected Clostridioides Infection (CDI) in Children</p>
Specimens required	<p>Send faecal specimens from any patient who has loose stools that score 5-7 on Bristol chart (appendix 1) and if no other cause of diarrhoea is known. If negative and loose stools persist, another two samples should be sent at 24-hour intervals. Relevant clinical information must be supplied with the specimen.</p> <p>There is no requirement to send clearance specimens from patients who become asymptomatic.</p> <p>Specimens should not be sent whilst patient is on treatment.</p> <p>Only when a relapse of CDI is suspected should you repeat the toxin testing and exclude other potential causes of diarrhoea, and only after 14 days of treatment. Relapse can also occur up to 14 days after therapy has stopped.</p>
Stool Charts	<p>It is the responsibility of staff looking after the patient within the area to record signs and symptoms of infection as appropriate, e.g. Bristol Stool Chart, Appendix 1. The date, time, size and nature of the stool should be recorded while symptomatic and continued until discharge in order to reduce</p>

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	15 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

	the risk of cross infection.
Surveillance	<p>Surveillance of CDI is mandatory in Scotland and is reported to HPS by the Diagnostic Laboratory.</p> <p>Local surveillance in NHSGGC is returned to wards with a prevalence of CDI monthly using Statistical Process Control Charts (SPCs). The trigger for action is when the numbers in a ward reach the upper control limit in the SPC. SPCs are not a substitute for local referral by clinical staff and ICTs but should be used to monitor trends and promote quality improvement.</p>
Terminal Cleaning of Room	<p>Follow SOP for Terminal Clean of Isolation Rooms. If isolation is discontinued and the patient remains in hospital, consider moving the patient to a new bed-space. This will allow the patient's bed, bed locker and bed table to be decontaminated thoroughly. These items can be expected, without cleaning, to remain contaminated.</p> <p>NB: relapse and re-infection from the environment can be as high as 20% in patients with CDI.</p> <p>See NHSGGC SOP Terminal Clean of Isolation Rooms</p>
Visitors	<p>Visitors are not required to wear aprons and gloves unless participating in patient care. If PPE is worn by patients or visitors it should be removed before leaving the room. Visitors should be advised to decontaminate their hands with liquid soap and water on leaving the room/ patient. Visitors should also be advised not to use communal areas or to sit on beds, while patient is infectious.</p>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	16 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

4. Evidence Base

Pai S et al. Five years experience of clostridium difficile infection in children at a UK tertiary hospital: proposed criteria for diagnosis and management. PLOS 2012; 71-6

Lees E A et al. The role of Clostridium difficile in the paediatric and neonatal gut — a narrative review. Eur J Clin Microbiol Infect Dis (2016) 35:1047–1057

<http://www.nipcm.hps.scot.nhs.uk/>

<https://www.hps.scot.nhs.uk/web-resources-container/guidance-on-prevention-and-control-of-clostridium-difficile-infection-cdi-in-health-and-social-care-settings-in-scotland/>

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	17 of 17
		Effective From	June 2020
	<i>Clostridioides difficile</i> Infection (CDI) in Children (3 -16 years) STANDARD OPERATING PROCEDURE	Review Date	June 2022
		Version	3
The most up-to-date version of this policy can be viewed at the following website: http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control			

Appendix 2 – Bowel Movement (adapted from the Bristol Stool Scale)

BOWEL MOVEMENT RECORD

Month: _____ Year: _____

Name: _____

Date	Time	Size S-small M-medium L-large S M L	Type 1	Type 2	Type 3	Type 4	Type 5	Type 6	Type 7	Staff Initials
			Separate hard lumps like nuts (hard to pass)	Sausage shaped but lumpy	Like a sausage but with cracks on surface	Like a sausage or snake, smooth and soft	Soft blobs with clear-cut edges (passed easily)	Fluffy pieces with ragged edges, a mushy stool	Watery, no solid pieces (entirely liquid)	
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									

Adapted from the Bristol Stool Scale developed by KW Heaton and SJ Lewis at the University of Bristol, 1997

The most up-to-date version of this policy can be viewed at the following website:
<http://www.nhsggc.org.uk/your-health/public-health/infection-prevention-and-control>